

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

THIS PAGE BLANK (USPTO)

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 231/22	A1	(11) International Publication Number: WO 98/34908 (43) International Publication Date: 13 August 1998 (13.08.98)
(21) International Application Number: PCT/IB97/00106 (22) International Filing Date: 11 February 1997 (11.02.97) (71) Applicant (for all designated States except US): BRACCO INTERNATIONAL B.V. [NL/NL]; Strawinskylaan 3051, NL-1077 ZX Amsterdam (NL). (72) Inventor; and (75) Inventor/Applicant (for US only): DESANTIS, Nicola [IT/IT]; Via Giosue Garducci, 14, I-20063 Cernusco sul Naviglio (IT). (74) Agent: MINOJA, Fabrizio; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).		(81) Designated States: AU, CA, CN, CZ, FI, HU, IL, JP, KR, MX, NO, NZ, PL, RU, UA, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PROCESS FOR THE CRYSTALLIZATION FROM A LINEAR OR BRANCHED (C5-C6) ALCOHOL OR THEIR MIXTURES OF (S)-N,N'-bis[2-HYDROXY-1- (HYDROXYMETHYL)ETHYL]-5-[(2-HYDROXY-1-OXOPROPYL)AMINO]-2,4,6- TRIIDO-1,3-BENZENDICARBOXAMIDE (57) Abstract Disclosed is a new process for the crystallisation of (S)-N,N'-bis[2-hydroxy-1- (hydroxymethyl)ethyl]-5-[(2-hydroxy-1-oxopropyl)amino] -2,4,6-triiodo-1,3-benzendicarboxamide (I), using a linear or branched (C5-C6) alcohol or their mixtures.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Process for the crystallization from a linear or branched (C5-C6) alcohol or their mixtures of (S)-N,N'-bis[2-hydroxy-1-(hydroxymethyl)ethyl]-5-[(2-hydroxy-1-oxopropyl)amino]-2,4,6-triiodo-1,3-benzendicarboxamide.

Technical Field

This invention refers to a new process for the crystallization of (S)-N,N'-bis[2-hydroxy-1-(hydroxymethyl)ethyl]-5-[(2-hydroxy-1-oxopropyl)amino]-2,4,6-triiodo-1,3-benzendicarboxamide (I), using a linear or branched (C5-C6) alcohol or their mixtures.

Background Art

The compound (I) is best known as Iopamidol, which is one of the world top compounds in the field of non-ionic X-ray contrast media.

The syntheses of Iopamidol known in literature, for instance the one described in GB 1472050, foresee a final purification at the end of the process, by using ion-exchange resins and successive recrystallization from EtOH or iso-PrOH.

Recently the patent application WO-A-95/04031 mentions different solvents (n-BuOH, sec-BuOH, iso-BuOH and/or t-BuOH) from which Iopamidol crystallizes. Butyl alcohols, especially 2-butanol, are used in a manner similar to that mentioned above for ethanol. It is claimed that butyl

alcohols are more easily removed than ethanol from the final product in the drying step.

A more recent application EP-A-0 747 344 discloses the use of propanol or isopropanol as crystallization solvent to overcome the problem of the formation of pasty products during the crystallization.

Another recent application IT95A001429 discloses the use of monoether of the ethylenglycol (e.g. Cellosolve® or Metilcellosolve®) as crystallization solvent for Iopamidol. This procedure allows the crystallization of Iopamidol starting from a product with a purity grade comprised between 99-99.5%.

Detailed Description of the Invention

We have now surprisingly found out that, and this is part of this invention, Iopamidol can be easily crystallized from the homologous (C5-C6) alcoholic solvents, with industrially acceptable yields and giving a product which meets the pharmacopoeia standards.

These solvents are very uncommon as crystallization solvent for their characteristics of high boiling point and low water solubility, especially for compound like Iopamidol very soluble in water. Therefore, the skilled technician would not have found a suggestion in the prior-art for using these alcohols.

The preferred solvents are selected from the following groups: hexanol, pentanol, 2-pentanol and 3-pentanol mixture, and iso-pentanol.

The improvement of this invention consists in the use of solvents able to give azeotropes with water forming two separate layers; the lower level has a minimal removable solvent content, while the upper one can be recycled.

This improvement is remarkable from the industrial point of view because it reduces the amount of solvent used and at the same time gives a product in accordance with pharmacopoeia standards, which have changed over the years (see the European Pharmacopoeia III ed. 1997 or USP XXIII-NF, 1996, V° suppl.) reducing the related substances content from 0.5% to 0.25%.

The achieved improvement, compared to the already used solvents cited in the above patents, is connected with the water percentage of the azeotropic mixture and the ability of the azeotrope to give two separate layers.

Despite WO-A-95/04031 describes the possibility to have the azeotropic mixture obtained by distillation separated in its components, that its true only for n-BuOH and iso-BuOH and not for all the claimed C4 alcohols, as cited in the following data regarding the azeotropes with water from "HANDBOOK OF CHEMISTRY AND PHYSICS", 68th Ed.

When the data are missing, experimental data are reported, obtained with the following procedure: 50 g of the solvent and 50 g of water are mixed in a reactor, equipped with stirrer, a Markusson settler. Then the mixture is refluxed,

the azeotropic layers are collected and the water content is determined by Karl-Fisher method.

SOLVENT	%WATER IN AZEOTROPE	%WATER UPPER LAYER	%WATER LOWER LAYER	AZEOTROPE TYPE
Ethanol	4.4%			homogeneous
1-Propanol	28.2%			homogeneous
iso-Propanol	12.2%			homogeneous
1-Butanol	44.5%	20.1%	92.3%	heterogeneous
2-Butanol	32%			homogeneous
iso-Butanol	30%	15.0%	91.3%	heterogeneous
tert-Butanol	11.75%			homogeneous
1-Pentanol	55%	12.4% (exp.)	95.05% (exp.)	heterogeneous
2-Pentanol	36.5%	12.8% (exp.)	86.2% (exp.)	heterogeneous
iso-Pentanol	49.6%	12.7% (exp.)	81.2% (exp.)	heterogeneous
3-Pentanol	35%	9.9%	94.5%	heterogeneous
n-Hexanol	67.2%	7.2%	99.42%	heterogeneous

It is easy to realize that the solvents of this invention, are those with a higher water percentage in the upper layer and form separate layers, except for n-BuOH and iso-BuOH. According to the process of this present invention, the recycled upper layer has a very low residual water content with respect to that of n-BuOH and iso-BuOH.

Furthermore, the residual water content present in the upper layer of the solvents of this invention is lower than 15% (w/w) and so that Iopamidol can be crystallized, if necessary, without anidrififying them.

Thus according to this invention the recycling of the organic layer becomes more advantageous from the industrial point of view since the costs for the solvents are curbed and the crystallization time are reduced.

Furthermore, as described in the Examples, the upper layer can be cooled and used it for washing the wet filtration cake. This procedure is absolutely new and is a great improvement from the industrial point of view.

The recovery of the solvent is easy and the upper layer can be directly used in the process of this invention without anidrififying it.

In this way the industrial process for the recovery of the solvent does not need elaborate systems or processes such as pervaporation or the simple addition of a little amount of a third solvent to the binary azeotropic mixture, for example toluene or cyclohexane, able to form a ternary azeotrope with water.

If the (C2-C4) alcohols are used, it is necessary to anidrifify the recovered solvent using, for example, one of the above cited procedures.

In the process of this invention a crude aqueous solution of Iopamidol 5-25% (w/w) is concentrated under vacuum at a pressure of 3-12 mmHg or atmospheric pressure, at a temperature comprised between 50 and 100°C to have a water

residual content comprised between 15-35% (w/w). Then the mixture is heated or cooled, depending on the case, and the crystallization solvent is added at a temperature comprised between 85-95°C, maintaining this temperature during the addition.

The amount of the solvent to be used is from 0.8 to 6 times (w/w) with respect to the amount of theoretical Iopamidol. Preferably, the solvent amount is from 0.8 to 4.5 times (w/w) with respect to the amount of theoretical Iopamidol. The water content is not considered, because water-saturated solvents can be used.

Then the mixture is azeotropic distilled, recycling the upper layer until the two layers are dissolved. Sometimes the distillation can be carried out to have a water residual content from 4 to 10% in the floating liquid. The solid can precipitate during this step, eventually by germination. In some cases it is possible to add the solvent in two or three consecutive portions.

In some cases, depending from the solvent, when the solid is beginning to crystallize in the lower layer, the distillation can be stopped and the mixture is cooled at the temperature from 60 to 80°C and then germinate.

After the precipitation of the solid, the distillation can be resumed until the water final content in the floating liquid is reached.

Then the temperature is taken to 17-25°C and maintained for 1-5 hours.

Then the solid is filtered, washed with dry solvents or with a quantity of the wet recycled solvent from the upper layer of the distilled azeotropic mixture.

The amount of the solvent is from 0.4 to 2 times (w/w) with respect to the amount of theoretical Iopamidol, preferably from 0.4 to 1 time.

The wet product is dried under a pressure comprised between 1 - 10 mmHg, preferably between 3 - 7 mmHg at a temperature comprised between 75°C and 95°C during at least 16 hours.

This procedure has no effect on the quality of the product and in this way the residual solvents are removed easily. The product is very stable, and decomposes at about 300°C without melting (see for example Merck Index 12th edition). After drying, no smell is detectable and surprisingly the residual content of the crystallization solvent is lower than 60 ppm.

The following table illustrates the content of the residual crystallization solvent in the final product and in the cited references, compared with the Examples of the present invention. In particular, the data regarding the ethanol as crystallization solvent are obtained using the procedure of the cited patent.

SOLVENT	RESIDUAL CONTENT	REFERENCES
Ethanol	490 ppm	GB 1472050
Ethanol	270 ppm	GB 1472050
Ethanol	370 ppm	GB 1472050

Sec-butanol	200 ppm	WO-A-9504031 (EX.1)
Sec-butanol	180 ppm	WO-A-9504031 (EX.2)
Sec-butanol	100 ppm	WO-A-9504031 (EX.3)
n-butanol	70 ppm	WO-A-9504031 (EX.4)
n-butanol	80 ppm	WO-A-9504031 (EX.5)
sec-butanol	100 ppm	WO-A-9504031 (EX.6)
t-butanol	150 ppm	WO-A-9504031 (EX.7)
n-pentanol	40 ppm	Example 1
iso-pentanol	30 ppm	Example 2
2-pentanol/3-pentanol	25 ppm	Example 3
n-hexanol	25 ppm	Example 4
n-hexanol	35 ppm	Example 5

Two preparations on industrial scale are reported in the Experimental part to demonstrate the improvement of the process of this invention.

All the reported examples give a product with a residual content of organic by-products lower than 0.25%, according to the more recent Pharmacopoeias.

The following examples serve to illustrate this invention and are not, in any way, to be considered as a limitation thereof. The water content in the azeotrope and in the final product was determined by the Karl-Fisher method, while the content of the solvent in the final product was determined by gas-chromatography.

EXAMPLE 1

1000 g of a 50% (w/w) crude aqueous solution of Iopamidol containing 0.4% of organic by-products is concentrated at atmospheric pressure at 100°C to obtain a water content comprised between 22% and 26% (w/w).

After cooling to about 95°C, 600 g of n-pentanol are added during an hour without cooling the solution. Then the solution is refluxed recycling the upper layer of the azeotrope mixture. The product precipitates during this step.

Then the mixture is distilled to have a 7-8% residue content of water in the floating liquid. Then the mixture is refluxed for 60 min. then cooled to 25°C and kept at this temperature for two hours. The product is filtered, washed with n-pentanol and dried at 80°C, 12 mmHg for 12 hours.

Yield: 94%

Residual solvent: 40 ppm

EXAMPLE 2

800 kg of a 70% (w/w) crude aqueous solution of Iopamidol containing 0.3% of organic by-products is concentrated at 12 mmHg pressure and 50°C to obtain a water content comprised between 25% and 30% (w/w) determined by Karl Fisher titration. Then the mixture is concentrated at atmospheric pressure at 100°C to obtain a residual water content comprised between 20% and 25% (w/w). 400 kg of iso-pentanol are added to the solution in 2 hours and very slowly in the first 15 min. The azeotrope is distilled using a distilling apparatus equipped with a settler recycling the upper phase. The distillation is continued to have a 10% (w/w) residual content of water determined by Karl Fisher titration. During this step the product precipitates spontaneously under crystalline form, without germination. Then 200 kg of dry iso-pentanol are added, maintaining the azeotropic distillation, to reach a 4-5% (w/w) of the residual content of water in the floating liquid. The mixture is refluxed for 1 hour, cooled in 3 hours to 15-20°C and finally filtered on centrifuge, washed with two portions of 100 kg of the recycled solvent from the upper phase of the azeotrope. Then the product is dried at 3 mmHg for 8 hours at 60°C and for other 8 hours at 75°C.

Yield: 95%

Residual solvent: 30 ppm

EXAMPLE 3

10000 g of a 50% (w/w) crude aqueous solution of Iopamidol containing 0.4% of organic by-products is concentrated at atmospheric pressure at 100°C to obtain a water content comprised between 22% and 26% (w/w). After cooling to about 95°C, 4000 g of a mixture of 30% of 2-pentanol and 70% of 3-pentanol in 3 hours is added without cooling the solution to 92°C. Then the solution is refluxed recycling the upper layer of the azeotrope mixture. The product precipitates during this step. Then the mixture is distilled to have a 5% residue content of water in the floating liquid. Then the mixture is refluxed for 60 min., then cooled to 25°C and kept at this temperature for 3 hours. The product is filtered, washed with 2000 g of the upper phase of the azeotrope mixture. Then the product is dried at 80°C with pressure of 12 mmHg during 12 hours.

Yield: 93%

Residual solvent: 25 ppm

EXAMPLE 4

100 g of a 70% (w/w) crude aqueous solution of Iopamidol containing 0.3% of organic by-products is concentrated at atmospheric pressure at 100°C to obtain a water content comprised between 22% and 35% (w/w). After cooling to about 95°C, 300 g of a n-hexanol saturated with water are added during about 1 hour without cooling the solution. Then the solution is refluxed recycling the upper layer of the azeotrope mixture. The product precipitates during this

step. Then the mixture is distilled to have a 7.2% residual water content in the floating liquid. Then the mixture is refluxed for 60 min., then cooled to 25°C and kept at this temperature for 2 hours. The product is filtered, washed with 50 g of the upper phase of the azeotrope. Then the product is dried at 80°C with pressure of 12 mmHg during 12 hours.

Yield: 90%

Residual solvent: 25 ppm

EXAMPLE 5

200 kg of a 70% (w/w) crude aqueous solution of Iopamidol containing 0.3% of organic by-products is concentrated at atmospheric pressure at 100°C to obtain a water content comprised between 22% and 35% (w/w). After cooling to about 95°C, 100 kg of a n-hexanol saturated with water are added during about 1 hour, without cooling the solution. Then the solution is refluxed recycling the upper layer of the azeotrope mixture to obtain the two layers system, where Iopamidol is beginning to crystallize in the lower layer. Then the mixture is cooled to a temperature comprised between 50-80°C and some germs are added to the mixture. Then the mixture is stirred at this temperature until the product terminates its crystallization (about 1 hour). Then the mixture is refluxed again recycling the upper layer of the azeotrope mixture distilled, to reach a 7.2% residual content of water in the floating liquid. The mixture is refluxed for 60 min., then cooled to 25°C and kept at this

temperature for 3 hours. The product is filtered, washed with two portions of 50 kg of the upper phase of the azeotrope.

The product is dried at 95°C with pressure of 3 mmHg during 20 hours.

Yield: 93%

Residual solvent: 35 ppm

EXAMPLE 6

200 kg of the mother liquors from the crystallization described in the Example 5, containing 7% of water, are distilled in a 300 L batch distillator, equipped with an impeller, a heating system using steam, to give a residue of about 50 kg. 130 kg of the upper layer with 7% of water and less than 1% of by-products (determined by GC) are recovered in the condenser and recycled.

CLAIMS

1. A process for the crystallization of Iopamidol which uses a linear or branched (C5-C6) alcohol or their mixtures.
2. A process according to claim 1, in which the crystallization solvent is selected in the group consisting of: hexanol, pentanol, 2-pentanol and 3-pentanol mixture, iso-pentanol.
3. A process according to claim 1, in which the crystallization solvent is hexanol.
4. A process according to claim 1, in which the crystallization solvent is pentanol.
5. A process according to claim 1, in which the crystallization solvent is 2-pentanol and 3-pentanol mixture.
6. A process according to claim 1, in which the crystallization solvent is iso-pentanol.
7. A process, according to claims 1-6, for the crystallization of Iopamidol to give said compound in a crystalline form and in accordance to the pharmacopeia standards comprising the following steps:

a) concentration under atmospheric or reduced pressure of an aqueous solution of Iopamidol 5-25% (w/w) at a temperature comprised between 50-100°C, to reach a residual water content comprised between 15-35%;

b) addition of the crystallization solvent at 85-95°C, maintaining this temperature during the addition;

c) distillation of the azeotropic mixture, recycling the upper layer until the two layers are dissolved; or termination of the distillation when the solid begins to crystallize in the mixture, cooling of the mixture at a temperature comprised between 60-80°C, and then germination;

d) the distillation can be resumed to reach a final water residual content comprised between 4-10% in the floating liquid;

e) the temperature is then taken to 17-25°C and kept for 1-5 hours;

f) filtration of the resulting precipitate, washed with the solvent;

g) under-vacuum drying of the solid under reduced pressure at a temperature comprised between 75-95°C, during at least 16 hours.

8. A process according to claim 7, in which the crystallization solvent is added in amounts from 0.8 to 6 times with respect to the amount of theoretical Iopamidol.

9. A process according to claim 7, in which the crystallization solvent is added in amounts from 0.8 to 4.5 times with respect to the amount of theoretical Iopamidol.

10. A process according to claim 7, in which the solvent for washing the filtrate is the wet recycled solvent from the upper layer of the azeotropic mixture cooled to room temperature.

11. A process according to claim 10, in which the solvent for washing the filtrate is added in amounts from 0.4 to 2 times with respect to the amount of theoretical Iopamidol.

12. A process according to claim 10, in which the solvent for washing the filtrate is added in amounts from 0.4 to 1 times with respect to the amount of theoretical Iopamidol.

13. Use of a linear or branched (C5-C6) alcohol or their mixtures, to crystallize Iopamidol.

14. Use of hexanol to crystallize Iopamidol.

15. Use of pentanol to crystallize Iopamidol.

16. Use of the 2-pentanol and 3-pentanol mixture to crystallize Iopamidol.

17. Use of iso-pentanol to crystallize Iopamidol.

18. Use of a linear or branched (C5-C6) alcohol or their mixtures, to obtain Iopamidol with characteristics corresponding to those required by pharmacopeia.

19. Use of hexanol to obtain Iopamidol with characteristics corresponding to those required by pharmacopeia.

20. Use of pentanol to obtain Iopamidol with characteristics corresponding to those required by pharmacopeia.

21. Use of 2-pentanol and 3-pentanol mixture to obtain Iopamidol with characteristics corresponding to those required by pharmacopeia.

22. Use of iso-pentanol to obtain Iopamidol with characteristics corresponding to those required by pharmacopeia.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/IB 97/00106

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07C231/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 04031 A (ZAMBON GROUP S.P.A., ITALY) 9 February 1995 cited in the application see claims	
A	--- EP 0 747 344 A (HOVIONE INTER LTD., SWITZ.) 11 December 1996 cited in the application see claims	1-22
A	--- WO 97 02235 A (RECORDATI S.A. CHEMICAL AND PHARMACEUTICAL COMPANY, SWITZ.; TARQUINI, A) 23 January 1997 cited in the application see claims	1-22
	--- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

- * "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- * "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- * "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * "&" document member of the same patent family

Date of the actual completion of the international search

18 September 1997

Date of mailing of the international search report

25.09.97

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Sánchez García, J.M.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 97/00106

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>GB 1 472 050 A (SAVAC AG) 27 April 1977 cited in the application see example 7</p> <p>-----</p>	1-22

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 97/00106

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9504031 A	09-02-95	IT 1257001 B AU 7532194 A BE 1008403 A CA 2128942 A CH 686783 A CN 1128020 A CZ 9401831 A DE 4426818 A DK 89294 A ES 2102312 A FI 960400 A FR 2708601 A GB 2280436 A,B HU 9600191 A IE 940609 A JP 7145125 A NL 9401229 A NO 960376 A NZ 271632 A PT 101554 A SE 9402595 A SI 9420046 A US 5571941 A ZA 9405472 A	05-01-96 28-02-95 07-05-96 31-01-95 28-06-96 31-07-96 15-02-95 02-02-95 31-01-95 16-07-97 29-01-96 10-02-95 01-02-95 28-11-96 08-02-95 06-06-95 16-02-95 29-01-96 27-07-97 04-05-95 31-01-95 31-10-96 05-11-96 03-03-95
EP 0747344 A	11-12-96	PT 101720 A AU 5451296 A CA 2178369 A CZ 9601632 A JP 8333319 A	31-01-97 19-12-96 09-12-96 11-12-96 17-12-96
WO 9702235 A	23-01-97	IT MI951429 A AU 6516196 A	07-01-97 05-02-97
GB 1472050 A	27-04-77	CH 608189 A AR 211853 A AT 349449 B AT 349450 B AT 340582 B AT 342579 B	29-12-78 31-03-78 10-04-79 10-04-79 27-12-77 10-04-78

INTERNATIONAL SEARCH REPORT

In relation to patent family members

International Application No.
PCT/IB 97/00106

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1472050 A		AU 475785 B	02-09-76
		AU 8751675 A	02-09-76
		BE 836355 A	08-06-76
		BG 26512 A	12-04-79
		CA 1046082 A	09-01-79
		CS 194234 B	30-11-79
		DE 2547789 A	24-06-76
		FR 2293919 A	09-07-76
		IN 142089 A	28-05-77
		JP 1097667 C	14-05-82
		JP 51082236 A	19-07-76
		JP 56042581 B	06-10-81
		NL 7512208 A,B	15-06-76
		SE 409993 B	17-09-79
		SE 7514076 A	14-06-76
		SI 7513155 A	31-12-94
		US 4001323 A	04-01-77
		ZA 7507222 A	24-11-76

THIS PAGE BLANK (USPTO)